

## KNEE PHYSICAL EXAM SMALL GROUP CASES

ACP Musculoskeletal Medicine Teaching Group  
ACP National Conference

1

### CASE #1: HISTORY

- Chief complaint: 35 year-old with bilateral knee pain.
- Onset: 2 years ago while in military service, hiking with heavy pack. No specific injury or event.
- Location: Anterior knee pain, around the patellae.
- Associated factors: No swelling. No locking or clicking. Reports painful knee "giving way" episodes, and feelings of painful instability when going downstairs\*. No falls.
- Exacerbating factors: Squatting, lunging, prolonged driving or sitting (theater sign), downhill/stairs worse than up. Gave up running because of pain.

2

### CASE #1: PHYSICAL EXAM

- Observation: Normal gait, no abnormality
- ROM: Full extension, full flexion but **pain anteriorly with terminal flexion.**
- Palpation: **Tenderness localized to the patellar facets,** otherwise no other tenderness. No effusion.
- Stability Testing: No laxity on anterior/posterior drawer, Lachman's, or varus/valgus stress testing.
- Provocative Testing: **Pain reproduced with squatting and patellar grind test.**

3

### PATELLOFEMORAL PAIN SYNDROME = ANTERIOR KNEE PAIN = RUNNER'S KNEE

Most common cause of knee pain in patients under 50 years old

- Definition
  - Diffuse, aching pain and stiffness in the anterior knee that increases with activities that place additional loads across the PF joint (squat, stairs, getting up from chair)
- Etiology
  - Multifactorial – malalignment, overuse, overloading
  - Degenerative changes = Patellofemoral OA (not PFPS)

Thomee R, et al. Patellofemoral pain syndrome: a review of current issues. Sports Med 1999; 28:245

Dixit S, et al. Management of patellofemoral pain syndrome. Am Fam Physician 2007; 75:194

Slide used with permission from Drs. Meg Pearson and Steve Bent

4

### PFPS: PHYSICAL EXAM

KEY: Diagnosis is based primarily on **history**

- Mechanism of injury: Rarely an acute injury, more often overuse or introduction of a new activity
- Pain: Behind or around patella, often bilateral. Mild to moderate
- Mechanical symptoms: Occasional popping sensation, “giving way”
- Exam
  - Often normal. Usually no atrophy or effusion
  - Patellar signs (facet tenderness and + grind test)
- Exacerbating factors – run, squat, stairs, sitting (theater sign).

Slide used with permission from Drs. Meg Pearson and Steve Bent

5

### PFPS: INITIAL PLAN

- Avoid exacerbating activities – squatting, lunging, running
- Find exercises that don’t exacerbate knee pain – often elliptical, cycling or swimming is tolerable
- Pain relief – Icing, NSAIDs, or Tylenol
- PT to optimize patellofemoral tracking – through strengthening of the core/hips, glutes, hamstrings, quads. Also, lower extremity stretching to correct flexibility issues.
- Weight-loss if indicated (10x force factor through PF joint)
- X-rays – usually not necessary. Only indicated initially if hx trauma or surgery, or presence of effusion. Consider in patients older than 50 to assess for PF OA

6

### PFPS: IF DOES NOT IMPROVE

- The prognosis is very good
- Referral to sports medicine physician if pain continues despite the above
- X-ray if not improving
- If x-rays unrevealing, consider MRI to evaluate the articular cartilage at the PF joint

7

### REFERENCES

- Kasitnon D, Li WX, Wang EXS, Fredericson M. [Physical Examination and Patellofemoral Pain Syndrome: an Updated Review](#). Curr Rev Musculoskelet Med. 2021 Dec;14(6):406-412. doi: 10.1007/s12178-021-09730-7. Epub 2021 Oct 29. PMID: 34713383
- Kuwabara A, Fredericson M. [Narrative Review of Anterior Knee Pain Differential Diagnosis \(Other than Patellofemoral Pain\)](#). Curr Rev Musculoskelet Med. 2021 Jun;14(3):232-238. doi: 10.1007/s12178-021-09704-9. Epub 2021 Apr 5. PMID: 33818700
- Collins NJ, Barton CJ, van Middelkoop M, Callaghan MJ, Rathleff MS, Vicenzino BT, Davis IS, Powers CM, Macri EM, Hart HF, de Oliveira Silva D, Crossley KM. [2018 Consensus statement on exercise therapy and physical interventions \(orthoses, taping and manual therapy\) to treat patellofemoral pain: recommendations from the 5th International Patellofemoral Pain Research Retreat, Gold Coast, Australia, 2017](#). Br J Sports Med. 2018 Sep; 52(18): 1170-1178
- Crossley KM, Stefanik JJ, Selfe J, Collins NJ, Davis IS, Powers CM, McConnell J, Vicenzino B, Bazett-Jones DM, Esculier JF, Morrissey D, Callaghan MJ. [2016 Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester. Part I: Terminology, definitions, clinical examination, natural history, patellofemoral osteoarthritis and patient-reported outcome measures](#). Br J Sports Med. 2016 Jul; 50(14):839-43

8

## CASE #2: HISTORY

- Chief complaint: 60 year-old slightly obese female with complaint of deep aching pain and stiffness in both knees.
- Onset: 3-year history of progressively worsening pain in both knees. Initially felt at work when lifting heavy materials, but more recently she has experienced pain in the absence of any physical activity or exertion.
- Location: medial knee pain bilaterally
- Associated factors: Intermittent swelling. Creaky, crackly sounds when knees move, not associated with increased pain. Occasionally feels as though one of her knees will give out. Denies knee catching or locking. Denies redness but sometimes experiences warmth or fullness in the knees.
- Exacerbating factors: Damp weather, kneeling, squatting, and stairs all worsen her pain. Her knees are stiff for about 20 minutes upon awakening and for a few minutes once getting up from a chair after prolonged sitting. She has difficulty walking > 30 minutes so no longer walks her dog.

9

## CASE #2: PHYSICAL EXAM

- Observation: **genu varum (bow-legged); mild antalgic gait; quadriceps atrophy.** No skin changes.
- ROM: **Passive ROM of both knees demonstrates palpable, painless crepitus. She is unable to flex or extend her knees completely. Maximum flexion is to about 120 degrees with pain.**
- Palpation: **Tenderness over medial joint lines and patellar facet bilaterally; right knee effusion**
- Stability Testing: No laxity on anterior/posterior drawer, Lachman's, or varus/valgus stress testing.
- Provocative Testing: (-)McMurray's test (painful but no pop/click). (-) Noble test.

10

## KNEE OSTEOARTHRITIS

- Metabolically active joint disease affecting:
  - Cartilage, bone, synovium/capsule and muscle
- Knee has 3 compartments – OA can affect one, some or all compartments
- Physical exam keys
  - Inspection: enlargement of bones, varus > valgus knees
  - Palpation: +/- effusion, lack of warmth, + crepitus, tender medial or lateral bony prominences (can also be tender at joint line)
  - ROM may be limited due to effusion or stiffness

11

## KNEE OA: INITIAL PLAN

- Avoid exacerbating activities – i.e. squatting, sitting for prolonged periods
- Weight loss – losing even a small amount of weight can significantly decrease knee pain from osteoarthritis (1 lb. wt. loss = 3-5 lbs. of decreased knee pressure)
- Exercise – strengthening the muscles around the knee makes the joint more stable and decreases pain. Stretching exercises help keep the knee joint mobile and flexible.
- Topical NSAIDS (diclofenac gel), oral NSAIDS, acetaminophen
- X-Ray – MUST be weight-bearing
- MRI not indicated in setting of OA without catching or locking symptoms.
- Consider intra-articular steroid injection
- Consider therapies – i.e. topical creams with capsaicin, acupuncture
- PT to strengthen muscles and increase flexibility
- Consider bracing if arthritis is in only one compartment
- Discuss referral to surgery when above measures are insufficient to allow maintenance of function.

12

### CASE #2 FURTHER HISTORY

After you explain the diagnosis of knee OA, the patient tells you that last week she actually had an MRI of the knees on the outside showing a medial meniscus tear. She brings in a copy of the report which shows a right degenerative medial meniscal tear. Does this change your management?

13

### CASE #3: HISTORY

- Chief complaint: 45 year-old with knee pain and swelling
- Onset: 4 weeks ago, walking and stepped off curb wrong and twisted knee
- Location: Medial joint line
- Associated factors: Swelling noted that evening, not immediately following injury. Swelling worse after activity. Clicking sensation with twisting. Denies instability or locking.
- Exacerbating factors: Squatting, twisting/pivoting on affected leg, walking long distances (swells).

14

## REFERENCES

- Faucett SC, Geisler BP, Chahla J, et al. Meniscus Root Repair vs Meniscectomy or Nonoperative Management to Prevent Knee Osteoarthritis After Medial Meniscus Root Tears: Clinical and Economic Effectiveness. *The American Journal of Sports Medicine*. 2019;47(3):762-769. doi:10.1177/0363546518755754
- Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the Hand, Hip, and Knee [published correction appears in *Arthritis Rheumatol*. 2021 May;73(5):799]. *Arthritis Rheumatol*. 2020;72(2):220-233. doi:10.1002/art.41142

15

## CASE #3: PHYSICAL EXAM

- Observation: **Mild effusion of affected knee.** No atrophy, skin changes, valgus/varus deformity.
- ROM: Full extension, full flexion but **pain over medial knee with deep knee flexion.**
- Palpation: **Tender over the medial joint line**, otherwise non-tender over patellar and quad tendons, lateral joint line, MCL/LCL, pes anserine bursa and patellar facets.
- Stability Testing: No laxity on Lachman's, varus/valgus stress testing or posterior drawer.
- Provocative Testing: **(+) McMurray's Test with pain medially. (+) Squat and Thessaly tests.**

16

## MENISCUS TEAR

- Mechanism of injury
  - Acute – twisting or squatting injury
  - Chronic – degenerative with minimal or no trauma
- Pain - joint line pain, sharp; may have popping, catching
- Effusion
  - 2-3 days after initial injury
  - Common after physical activity
- Exacerbating factors
  - Twisting, squatting, deep knee flexion

Slide used with permission from Drs. Meg Pearson and Steve Bent

17

## MENISCUS TEAR PHYSICAL EXAM

- Inspection: might notice effusion
- Palpation: small – moderate effusion
- ROM: effusion or bucket handle tear can limit ROM
- Palpation: isolated joint line tenderness of affected side
- Provocative Tests
  - Squat
  - McMurray
  - Thessaly

18

### MENISCUS TEAR: INITIAL PLAN

- Avoid exacerbating activities – squatting, twisting
- Continue or change NSAID, consider APAP
- May obtain imaging at first visit as patient has failed 4 weeks of home conservative care and continues to have mechanical clicking symptoms and pain/swelling
- -X-ray – MUST be weight-bearing: Normal as above
- -MRI – only AFTER x-ray is obtained to rule out any OA: meniscus tear as above
- Referral to Orthopedics for consideration of knee arthroscopy with partial meniscectomy vs meniscus repair
- PT for LE conditioning prior to surgery if pt. also has OA
- Obtain urgent MRI and refer urgently to Orthopedics if bucket handle meniscus tear or meniscal root tear

19

### CASE #3B

- WOULD YOU CHANGE YOUR MANAGEMENT IF THE X-RAY SHOWED:

*Bilateral mild/moderate joint space narrowing, worse medially. Subchondral sclerosis, peaking of tibial spines, and several osteophytes.*

- WHAT WOULD YOU DO DIFFERENTLY?

20

## REFERENCES

- Abram SGF, Hopewell S, Monk AP, Bayliss LE, Beard DJ, Price AJ. Arthroscopic partial meniscectomy for meniscal tears of the knee: a systematic review and meta-analysis. *Br J Sports Med.* 2020 Jun;54(11):652-663. doi: 10.1136/bjsports-2018-100223. Epub 2019 Feb 22. PMID: 30796103.
- van de Graaf VA, Noorduyn JCA, Willigenburg NW, Butter IK, de Gast A, Mol BW, Saris DBF, Twisk JWR, Poolman RW; ESCAPE Research Group. Effect of Early Surgery vs Physical Therapy on Knee Function Among Patients With Nonobstructive Meniscal Tears: The ESCAPE Randomized Clinical Trial. *JAMA.* 2018 Oct 2;320(13):1328-1337. doi: 10.1001/jama.2018.13308. Erratum in: *JAMA.* 2018 Dec 4;320(21):2272-2273. Erratum in: *JAMA.* 2020 Mar 10;323(10):1001. PMID: 30285177; PMCID: PMC6583004.
- Lee SH, Lee OS, Kim ST, Lee YS. Revisiting Arthroscopic Partial Meniscectomy for Degenerative Tears in Knees With Mild or No Osteoarthritis: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Clin J Sport Med.* 2020 May;30(3):195-202. doi: 10.1097/JSM.0000000000000585. PMID: 29995671.
- Sihvonen R, Paavola M, Malmivaara A, Itälä A, Joukainen A, Kalske J, Nurmi H, Kumm J, Sillanpää N, Kiekara T, Turkiewicz A, Toivonen P, Englund M, Taimela S, Järvinen TLN; FIDELITY (Finnish Degenerative Meniscus Lesion Study) Investigators. Arthroscopic partial meniscectomy for a degenerative meniscus tear: a 5 year follow-up of the placebo-surgery controlled FIDELITY (Finnish Degenerative Meniscus Lesion Study) trial. *Br J Sports Med.* 2020 Nov;54(22):1332-1339. doi: 10.1136/bjsports-2020-102813. Epub 2020 Aug 27. PMID: 32855201; PMCID: PMC7606577.
- Faucett SC, Geisler BP, Chahla J, et al. Meniscus Root Repair vs Meniscectomy or Nonoperative Management to Prevent Knee Osteoarthritis After Medial Meniscus Root Tears: Clinical and Economic Effectiveness. *The American Journal of Sports Medicine.* 2019;47(3):762-769. doi:10.1177/0363546518755754

21

## CASE #4: HISTORY

- Chief complaint: 47 year-old with right knee pain
- Onset: 2 weeks ago after a fall on a paddle board. His foot slipped out from under him and he felt a pop and acute pain at the medial knee when his knee fell into valgus
- Location: medial joint line
- Associated factors: no swelling, clicking or catching. Denies locking. **Reports that his knee “feels loose,” particularly when changing direction or turning to his left.**

22

#### CASE #4: PHYSICAL EXAM

- Observation: **abnormal, antalgic gait, cautious.** He prefers to sit with his hip in external rotation with weight on the lateral foot and a varus load on the knee
- ROM: full ROM, painful medially with full flexion
- Effusion: none
- Palpation: **point tenderness at the MCL, both proximal and distal to the medial joint line.** Not tender elsewhere on the joint lines or other parts of the knee
- Stability Testing: **(+) Laxity on valgus stress at 30 degrees;** No laxity with valgus stress at 0 degrees. No laxity on Lachman's, varus stress testing or posterior drawer
- Provocative Testing: (-) McMurray's test.

23

#### MCL TEAR

- History
  - Force hits outside of the knee causing MCL to stretch and possibly tear (valgus stress to knee)
  - Pain medial knee
  - Knee feels stiff, pain with terminal flexion due to tension on ligament
  - Mild swelling
  - +/- instability

24

**MCL TEAR:  
SEVERITY GRADED BY DEGREE OF LAXITY WITH  
VALGUS STRESS AT 30 DEGREES**

Grade	Injury	Translation compared to unaffected side	Patient response
I	Strain	Minimal laxity, firm endpoint	Pain
II	Partial tear	Some laxity, firm endpoint	Pain, may feel loose
III	Complete tear	Obvious laxity, no endpoint	Minimal pain, may feel very loose

25

**MCL SPRAIN, GRADE 2: INITIAL PLAN**

- Double-hinged knee brace, restricted ROM (i.e. 0-80 degrees) via brace for 3 weeks
- PT referral
- NSAIDs, ice – to control pain and decrease inflammation
- Avoid exacerbating activities

26

## REFERENCES

- Miyamoto, Ryan G. MD; Bosco, Joseph A. MD; Sherman, Orrin H. MD Treatment of Medial Collateral Ligament Injuries, Journal of the American Academy of Orthopaedic Surgeons: March 2009 - Volume 17 - Issue 3 - p 152-161

27

## CASE #5: HISTORY

- Chief complaint: 35 year-old woman with acute knee pain during trampoline half-pipe.
- Onset: 3 days ago jumped down and felt a pop with immediate knee pain and swelling. Went to ER: placed in knee immobilizer and given NSAIDS.
- Location: Posterior and lateral knee pain with terminal extension, tightness with bending.
- Associated factors: Knee feels unstable if not in the brace, especially with cutting or pivoting.
- Exacerbating factors: Instability with squatting, pivoting, walking down stairs, and stepping laterally (entire body weight on affected leg)

28

### CASE #5: PHYSICAL EXAM

- Observation: **Using crutches, leg in immobilizer, large knee effusion.** No skin changes, no valgus/varus deformity.
- ROM: **Lacks 5 degrees of extension, can only flex the knee to 100 degrees, limited due to pain** (*teaching point: determine if knee is locking—suggests concomitant meniscal tear- or if ROM is limited due to effusion*)
- Palpation: **tender to palpation on the lateral femoral condyle and lateral tibial plateau**
- Stability Testing: **(+) Lachman's, +Anterior drawer.**
- Provocative Testing: **(-) McMurrays. Deferred Thessaly and squat 2/2 instability**

29

### ANTERIOR CRUCIATE LIGAMENT (ACL) TEAR

- Main stabilizing ligament of the knee
- MOI
  - Twisting injury to knee (non-contact)
- Rate is higher in women in certain sports



30

## ACL TEAR

- History
  - Often feel a pop
  - Swelling develops soon after the injury
  - May be unstable upon pivoting
- PE
  - Effusion
  - Tenderness
    - +/- medial or lateral
  - +Lachman test
    - Very high sensitivity (0.87) and specificity (0.98)

31

## ACL TEAR: INITIAL PLAN

- Rest, ice, compression, elevation
- X-rays (to quickly evaluate for a fracture), MRI (has a sensitivity of 0.97 and specificity of 1.0 for detecting ACL injuries)
- Discontinue knee immobilizer to prevent muscle atrophy, but consider double-hinged knee brace.
- If knee is unstable, crutches to avoid weight-bearing if intolerant of double-hinged brace
- OTC analgesics sufficient for most cases: Tylenol and NSAIDS
- Consider aspiration +/- steroid injection to relieve effusion/hemarthrosis and help with pre-op goals of 1) walking without a limp, 2) return of full extension ROM, and 3) no effusion
- Operative vs non-op management:
  - Most active, younger patients and high-level athletes opt for surgery.
  - For older patients and/or poor surgical candidates, trial non-op is reasonable
  - Ortho referral for most patients who are surgical candidates to discuss options.
  - Patients who decide not to pursue surgical management should be referred to PT or primary care sports medicine

32

## REFERENCES

- Musahl V, Karlsson J. Anterior Cruciate Ligament Tear. N Engl J Med. 2019 Jun 13;380(24):2341-2348. doi: 10.1056/NEJMc1805931. PMID: 31189037.
- Filbay SR, Grindem H. Evidence-based recommendations for the management of anterior cruciate ligament (ACL) rupture. Best Pract Res Clin Rheumatol. 2019;33(1):33-47. doi:10.1016/j.berh.2019.01.018

33

## CASE #6: HISTORY

- Chief complaint: 27 year-old presents with right knee pain
- Onset: 2 weeks ago while training for his second sprint triathlon. Has been increasing training volume rapidly.
- Location: just above the lateral joint line
- Associated factors: no swelling, clicking or catching. Denies instability or locking.
- Exacerbating factors: climbing stairs, running downhill or long distance. Cycling makes the pain worse. Pain resolves shortly after running and cycling.

34

### CASE #6: PHYSICAL EXAM

- Observation: **abnormal gait (walking with the affected knee extended)**
- ROM: full extension and flexion.
- Palpation: **point tenderness upon palpation of the lateral femoral condyle and at the lateral tubercle of the tibia (a.k.a Gerdy's tubercle).**
- Stability Testing: No laxity on Lachman's, varus/valgus stress or posterior drawer testing.
- Provocative Testing: **(+)Noble test.** (-) McMurray's, squat and Thessaly tests.

35

### IT BAND FRICTION SYNDROME

- History
  - Pain at lateral femoral condyle
  - Due to repetitive flexion/extension motion of knee (i.e running, cycling)
  - As ITB becomes increasingly irritated, symptoms typically begin earlier in an exercise session
- PE
  - TTP of lateral knee approx. 2 cm above j line
  - TTP worse when patient is standing and knee is flexed to 30 degrees
  - Positive Noble, Ober's



36

### ITB SYNDROME: INITIAL PLAN

- Avoid exacerbating activities – i.e. running on pavement, cycling. Consider low impact activities (swimming) to maintain cardiovascular fitness until inflammation reduced
- PT referral to correct mobility and strength deficits
- Topical or oral NSAIDs, ice
- Refresh running shoes
- Identify anatomic factors that may contribute to ITBS:
  - Leg-length discrepancy: consider a prescription heel-lift (to correct 50% of the leg length discrepancy, can do 100% if 50% doesn't help)

37

### ITB SYNDROME: FOLLOW-UP PLAN

- After acute pain has resolved, start hip/core strengthening and a stretching program
- Avoid running on banked surfaces and/or hills or running in the same direction on a track.
- Cyclists: Suggest a bike fit
- If not improving consider referral to sports medicine or ortho

38

## REFERENCES

- Khaund R, Flynn S. IT Band Syndrome. Am Fam Physician 2005;71:1545-50.
- Beals C, Flanigan D. Review of Treatments for ITBS in Athletic Population. J of Sports Med Vol 2013. <http://dx.doi.org/10.1155/2013/367169>
- Geisler PR. Current Clinical Concepts: Synthesizing the Available Evidence for Improved Clinical Outcomes in Iliotibial Band Impingement Syndrome. J Athl Train. 2021 Aug 1;56(8):805-815. doi: 10.4085/1062-6050-548-19. PMID: 34375405; PMCID: PMC8359713.
- Jackson J. Iliotibial band syndrome. UpToDate. Literature review current through Dec 2022, topic last updated May 20, 2022. Accessed January 30, 2023.